arXiv:1001.3095v1 [math.DG] 18 Jan 2010

Dual leftinvariant almost complex structures on the $SU(2) \times SU(2)$.

Natalia Daurtseva
Russia, Kemerovo State University
natali0112@ngs.ru

Key-words: nearly Kähler structure, leftinvariant 3-form, almost complex structure

Abstract

There exist non-degenerate 3-form $d\omega_I$, $\omega_I(X,Y) = g(IX,Y)$, for each leftinvariant almost Hermitian structure (g,I), where g is Killing-Cartan metric on the $M = S^3 \times S^3 = SU(2) \times SU(2)$. Known [3], that arbitrary non-degenerate 3-form on the 6-dimensional manifold, with some additional properties defines the almost complex structure. Condition for I to define almost complex structure J_I by $d\omega_I$ is obtained. Properties of J_I are researched.

Introduction. Let M^6 be a smooth (i.e. C^∞), closed, orientable manifold, with differential 3-form $\psi \in \Lambda^3(M)$. The form $\psi \in \Lambda^3(M)$ defines [3] endomorphism $K \in \mathrm{End}(TM) \otimes \Lambda^6(M)$ by

$$K(X) = A(i_X\psi \wedge \psi),$$

where $A : \Lambda^5 \longrightarrow TM \otimes \Lambda^6$ is the isomorphism induced by the exterior product ($i_{A(\varphi)}Vol = \varphi$, where Vol is fixed volume form, $A(\varphi)$ is the suitable vector field here). Let denote $\frac{1}{6}\mathrm{tr}K^2$ as $\tau(\psi)$, then $K^2 = \mathrm{Id} \otimes \tau(\psi)$.

The group $GL(6,\mathbb{R})$ has two open orbits $\mathcal{O}_1$ and $\mathcal{O}_2$ on $\Lambda^3(\mathbb{R}^6)$. The stabilizer of the forms in the first orbit is $SL(3,\mathbb{C})$. Known [4], that $\tau(\psi) < 0$ if and only if $\psi \in \mathcal{O}_1$. Thus, if differential 3-form ψ belongs to $\mathcal{O}_1$ at each point, then it determines the almost complex structure J on M:

$$J = \frac{1}{\kappa}K,$$

where $\kappa = \sqrt{-\tau(\psi)}$.

Suppose, that now two forms (ω, ψ), $\omega \in \Lambda^2(M)$ $(\omega \wedge \omega \wedge \omega \neq 0)$, $\psi \in \Lambda^3(M)$ are defined on the M. The 3-form ψ, with $\tau(\psi) < 0$ provides a reduction to $SL(3,\mathbb{C})$ and 2-form ω to $Sp(3,\mathbb{R})$. As the group $SU(3) = Sp(3,\mathbb{R}) \bigcap SL(3,\mathbb{C})$, then two forms (ω, ψ) with some compatibility conditions provide a reduction to $SU(3)$.

The first of this conditions is:

$$\omega \wedge \psi = 0$$

This says that ω is of type (1,1) with respect to the above almost complex structure J. The second condition is that $\omega(X, JX)$ has to be positive definite form. It gives the Hermitian structure (g, ω, J) on M, where $g(X,Y) = \omega(X, JY)$.

Further, if pair (ω, ψ) gives a reduction to $SU(3)$ and:

$$\begin{cases} \psi = 3d\omega; \\ d\phi = -2\mu\omega \wedge \omega, \text{ where } i_X\psi = i_{JX}\phi, \ \mu \in \mathbb{R} \end{cases}$$

then $(M, g, J) \in \mathcal{NK}$ [2]. This approach is used in [2] to construct invariant nearly Kähler structure on $S^3 \times S^3$.

Let $M = S^3 \times S^3$ now. We'll take interest in leftinvariant structures on $S^3 \times S^3 = SU(2) \times SU(2)$. In this case all the calculations are reduced to ones on the Lie algebra $\mathfrak{su}(2) \times \mathfrak{su}(2)$ of Lie group $SU(2) \times SU(2)$. Denote $(e_1, e_2, e_3, e_4, e_5, e_6)$ the standard frame of $\mathfrak{su}(2) \times \mathfrak{su}(2) = \mathbb{R}^3 \times \mathbb{R}^3$ ($[e_1, e_2] = e_3$, $[e_1, e_3] = -e_2$, $[e_2, e_3] = e_1$, $[e_4, e_5] = e_6$, $[e_4, e_6] = -e_5$, $[e_5, e_6] = e_4$, $[e_i, e_j] = 0$, for other i, j; $e_i \in \mathfrak{su}(2) \times \{0\}$, $i = 1, 2, 3$; $e_i \in \{0\} \times \mathfrak{su}(2)$, $i = 4, 5, 6$). Let fix the orientation on M, which defined by the selection of vectors $(e_1, e_2, e_3, e_4, e_5, e_6)$. Consider the space $\mathcal{A}^+$ of all leftinvariant almost complex structures on M that induce the given orientation.

Fix the Riemannian metric g induced by the Killing-Cartan form on $SU(2) \times SU(2)$. Take the space $\mathcal{AO}_g^+ \subset \mathcal{A}^+$ of all leftinvariant g-orthogonal almost complex structures J, induced the given orientation on $SU(2)\times SU(2)$. Known [6], that $\mathcal{A}^+$ is bundle over the $\mathcal{AO}_g^+$, fiber of the bundle over the $J \in \mathcal{AO}_g^+$ is the space $\mathcal{A}^+_{\omega_J} = \{I \in \mathcal{A}^+ : \ \omega_J(IX, IY) = \omega_J(X, Y), \ \omega_J(X, IX) > 0, \forall X, Y \in \mathfrak{su}(2) \times \mathfrak{su}(2)\}$ of almost complex structures, positive associated with non-degenerate form $\omega_J(X,Y) = g(JX, Y)$.

The leftinvariant 2-form $\omega_I(X,Y) = g(IX, Y)$ corresponds to each almost complex structure $I \in \mathcal{AO}_g^+$. Known that closed non-degenerate differential 2-form does not exist on $SU(2) \times SU(2)$. Thus $d\omega_I \neq 0$. Therefore the map:

$$I \in \mathcal{AO}_g^+ \longrightarrow d\omega_I \longrightarrow K \in \mathrm{End}(\mathfrak{su}(2) \times \mathfrak{su}(2))$$

is defined. If $\tau(d\omega_I) < 0$, then $I \in \mathcal{AO}_g^+$ defines almost complex structure $J_I \in \mathcal{A}^+$.

The almost complex structures $I \in \mathcal{AO}_g^+$, for which $\tau(d\omega_I) < 0$ are investigated in the paper. The properties of corresponding almost complex structures $J_I \in \mathcal{A}^+$ are studied.

Calculation. Let $I \in \mathcal{AO}_g^+$. It is defined by skew-symmetric matrix

$$I = \begin{pmatrix} A & B \\ -B^T & C \end{pmatrix},$$

where $A = \begin{pmatrix} 0 & a_1 & a_2 \\ -a_1 & 0 & a_3 \\ -a_2 & -a_3 & 0 \end{pmatrix}, B = \begin{pmatrix} b_1 & b_2 & b_3 \\ b_4 & b_5 & b_6 \\ b_7 & b_8 & b_9 \end{pmatrix}, C = \begin{pmatrix} 0 & c_1 & c_2 \\ -c_1 & 0 & c_3 \\ -c_2 & -c_3 & 0 \end{pmatrix}.$ in the standard frame. Parameters $a_i, b_j, c_k,\ i,k = 1,2,3,\ j = 1,..,6$ are in relations given by condition $I^2 = -1$, in particular we have:

$$\begin{aligned}
&1)\ b_1^2 + b_2^2 + b_3^2 + a_1^2 + a_2^2 = 1. \\
&2)\ b_4^2 + b_5^2 + b_6^2 + a_1^2 + a_3^2 = 1. \\
&3)\ b_7^2 + b_8^2 + b_9^2 + a_2^2 + a_3^2 = 1. \\
&4)\ a_2a_3 + b_1b_4 + b_2b_5 + b_3b_6 = 0. \\
&5)\ -a_1a_3 + b_1b_7 + b_2b_8 + b_3b_9 = 0. \\
&6)\ a_1a_2 + b_4b_7 + b_5b_8 + b_6b_9 = 0. \\
&7)\ b_1^2 + b_4^2 + b_7^2 + c_1^2 + c_2^2 = 1. \\
&8)\ b_2^2 + b_5^2 + b_8^2 + c_1^2 + c_3^2 = 1. \\
&9)\ b_3^2 + b_6^2 + b_9^2 + c_2^2 + c_3^2 = 1.
\end{aligned}$$

Theorem 1. $\tau(d\omega_I) < 0$ *if and only if* $a_1^2 + a_2^2 + a_3^2 < \frac{3}{4}$.

Proof. Let use the Maurer-Cartan formulas to calculate $d\omega_I$: $d\theta^k = -\sum_{i<j} C_{ij}^k \theta^i \wedge \theta^j$, where $C_{ij}^k,\ k,i,j = 1\dots6$ are the structural constants, $\{\theta^i, i = 1\dots6\}$ the frame of the leftinvariant 1-forms space on $M = SU(2) \times SU(2)$. Let $(e^1, e^2, e^3, e^4, e^5, e^6)$ is co-frame to $(e_1, e_2, e_3, e_4, e_5, e_6)$. Then $de^1 = -e^2 \wedge e^3$, $de^2 = -e^3 \wedge e^1$, $de^3 = -e^1 \wedge e^2$, $de^4 = -e^5 \wedge e^6$, $de^5 = -e^6 \wedge e^4$, $de^6 = -e^4 \wedge e^5$. We have

$$\psi = d\omega_I = b_1(e^{234} - e^{156}) + b_2(e^{235} - e^{164}) + b_3(e^{236} - e^{145}) + b_4(e^{314} - e^{256}) +$$

$$+ b_5(e^{315} - e^{264}) + b_6(e^{316} - e^{245}) + b_7(e^{124} - e^{356}) + b_8(e^{125} - e^{364}) + b_9(e^{126} - e^{345})$$

where e^{ijk} is the 3-form $e^i \wedge e^j \wedge e^k$.

$$i_{e_1}\psi \wedge \psi = (-b_1^2 - b_2^2 - b_3^2 + b_4^2 + b_5^2 + b_6^2 + b_7^2 + b_8^2 + b_9^2)e^{23456} +$$

$$+ 2(b_1b_4 + b_2b_5 + b_3b_6)e^{13456} - 2(b_1b_7 + b_2b_8 + b_3b_9)e^{12456} +$$

$$+2(b_6b_8-b_5b_9)e^{12356}+2(b_6b_7-b_4b_9)e^{12346}+2(b_5b_7-b_4b_8)e^{12345}$$

Then

$$i_{e_1}\psi\wedge\psi=(1-2a_3^2)e^{23456}-2a_2a_3e^{13456}-2a_1a_3e^{12456}+2(b_6b_8-b_5b_9)e^{12356}+$$

$$+2(b_6b_7-b_4b_9)e^{12346}+2(b_5b_7-b_4b_8)e^{12345}$$

As $i_{K(X)}Vol=i_X\psi\wedge\psi$, and $i_YVol=Y^1e^{23456}-Y^2e^{13456}+Y^3e^{12456}-Y^4e^{12356}+Y^5e^{12346}-Y^6e^{12345}$, where $Vol=e^1\wedge e^2\wedge e^3\wedge e^4\wedge e^5\wedge e^6$, then matrix of $\frac{1}{2}K$ in the above basis is

$$\begin{pmatrix}
\frac{1}{2}-a_3^2 & a_2a_3 & -a_1a_3 & b_8b_6-b_9b_5 & b_9b_4-b_7b_6 & b_7b_5-b_8b_4\\
a_2a_3 & \frac{1}{2}-a_2^2 & a_1a_2 & b_9b_2-b_8b_3 & b_7b_3-b_1b_9 & b_8b_1-b_7b_2\\
-a_1a_3 & a_1a_2 & \frac{1}{2}-a_1^2 & b_5b_3-b_6b_2 & b_6b_1-b_4b_3 & b_4b_2-b_5b_1\\
b_9b_5-b_8b_6 & b_8b_3-b_9b_2 & b_2b_6-b_3b_5 & -\frac{1}{2}+c_3^2 & -c_2c_3 & c_1c_3\\
b_7b_6-b_9b_4 & b_1b_9-b_3b_7 & b_4b_3-b_6b_1 & -c_2c_3 & -\frac{1}{2}+c_2^2 & -c_1c_2\\
b_8b_4-b_7b_5 & b_2b_7-b_8b_1 & b_5b_1-b_4b_2 & c_1c_3 & -c_1c_2 & -\frac{1}{2}+c_1^2
\end{pmatrix}$$

We can calculate $(K^2)_{11}$:

$$(K^2)_{11}=(1-2a_3^2)^2+4a_2^2a_3^2+4a_1^2a_3^2-4(b_8b_6-b_9b_5)^2-4(b_9b_4-b_7b_6)^2-4(b_7b_5-b_8b_4)^2$$

Denote

$$x=a_1^2+a_2^2+a_3^2$$

Then

$$(K^2)_{11}=1-4a_3^2+4a_3^2x-4b_8^2b_6^2-4b_9^2b_5^2+8b_8b_6b_9b_5-4b_9^2b_4^2-4b_7^2b_6^2+8b_9b_4b_7b_6-$$

$$-4b_7^2b_5^2-4b_8^2b_4^2+8b_7b_5b_8b_4=1-4a_3^2(1-x)-4b_4^2(b_7^2+b_8^2+b_9^2)-4b_5^2(b_7^2+b_8^2+b_9^2)-$$

$$-4b_6^2(b_7^2+b_8^2+b_9^2)+4b_4^2b_7^2+4b_5^2b_8^2+4b_6^2b_9^2+8b_8b_6b_9b_5+8b_9b_4b_7b_6+8b_7b_5b_8b_4=$$

$$=1-4a_3^2(1-x)-4(b_4^2+b_5^2+b_6^2)(b_7^2+b_8^2+b_9^2)+4(b_4b_7+b_5b_8+b_6b_9)^2$$

Use the condition $I^2=-1$, obtain:

$$(K^2)_{11}=1-4a_3^2(1-x)-4(1-a_1^2-a_3^2)(1-a_2^2-a_3^2)+4(-a_1a_2)^2=4x-3$$

As $(K^2)_{11}=(K^2)_{ii}$, $\forall i=1..6$, we have

$$\tau(d\omega_I)=\frac{1}{6}\mathrm{tr}K^2=-3+4(a_1^2+a_2^2+a_3^2)$$

□

Remark 1. One can write the matrix of K in the frame (e):

$$K = \begin{pmatrix} 1 - 2A^* & -2B^* \\ 2B^{*T} & -1 + 2C^* \end{pmatrix},$$

where A^*, B^*, C^* - are the algebraic supplement matrices to A, B, C respectively.

Remark 2. As $a_1^2 + a_2^2 + a_3^2 = c_1^2 + c_2^2 + c_3^2$ and $\sum_{i=1}^{9} b_i^2 = 3 - 2\sum_{i=1}^{3} a_i^2$ by $I^2 = -1$, then the conditions:
1. $\tau(d\omega_I) < 0$,
2. $c_1^2 + c_2^2 + c_3^2 < \frac{3}{4}$,
3. $a_1^2 + a_2^2 + a_3^2 < \frac{3}{4}$,
4. $\sum_{i=1}^{9} b_i^2 > \frac{3}{2}$

are equivalent.

Denote the set of all $I \in \mathcal{AO}_g^+$, with $\tau(d\omega_I) < 0$ as $\mathcal{AO}_-$. Now we can define map $\alpha : \mathcal{AO}_- \longrightarrow \mathcal{A}^+$, $\alpha(I) = J_I = \frac{1}{\sqrt{-\tau(d\omega_I)}}K$ for each $I \in \mathcal{AO}_-$.

Remark 3. $\tau(d\omega_J) = 1 > 0$ for integrable almost complex structure $J = \begin{pmatrix} A & B \\ -B & A \end{pmatrix}$, $A = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}$, $B = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$. For $I_0 = \begin{pmatrix} 0 & -E \\ E & 0 \end{pmatrix}$ we have $\tau(d\omega_I) = -3$, then $J_{I_0} = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & -2 \\ 2 & -1 \end{pmatrix}$ with metric $g_{\mathcal{NK}} = \frac{1}{\sqrt{3}} \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$ gives nearly Kähler structure.

Lemma 1. $\det B < 0$ *for all almost complex structures* $I \in \mathcal{AO}_-$.

Proof. Assume that the almost complex structure $I \in \mathcal{AO}_-$, such that $\det B = 0$ exists. Then we can find non-zero vector $v \in \mathbb{R}^3$, for which $B^T v = 0$. Let $V \in \mathfrak{su}(2) \times \mathfrak{su}(2)$, such that its projection on the first factor is v $(\pi_1(V) = v)$ and its projection on the second factor is zero $(\pi_2(V) = 0)$, i.e. $V^T = (v^T, 0)$. Then

$$(I^2 V)^T = (v^T A^2, v^T AB) = (-v^T, 0) \;\Leftrightarrow\; \begin{cases} A^2 v = -v; \\ B^T A v = 0. \end{cases}$$

Condition $A^2 v = -v$ is equivalent to $\det(A^2 + 1) = 0$, i.e. $x = 1$, this one contradicts to $x < 3/4$.

As $\det B \neq 0$, then vectors (e_1, e_2, e_3) are I-linearly non-dependent. Then

$$(e_1, e_2, e_3, Ie_1, Ie_2, Ie_3) = (e_1, e_2, e_3, e_4, e_5, e_6) \begin{pmatrix} E & A \\ 0 & -B^T \end{pmatrix}$$

I.e. I keeps the orientation [5] if $\det B < 0$.

$\square$

Lemma 2. $J_I \in \mathcal{A}^+$.

Proof. As $\det B \neq 0$, then (e_1, e_2, e_3) are J_I-linearly non-dependent. The determinant of amplication matrix from the frame $(e_1, e_2, e_3, J_Ie_1, J_Ie_2, J_Ie_3)$ to the standard one is

$$\det \begin{pmatrix} E & 1-2A^* \\ 0 & 2B^{*T} \end{pmatrix} = 8(\det B)^2 > 0.$$

$\square$

Theorem 2. *$J_I \in \mathcal{A}^+_{\omega_J}$, where*

$$\omega_J = \frac{2}{\sqrt(1-\tau)} \begin{pmatrix} 0 & (1+yA^*)B^* \\ -(1+yC^*)B^{*T} & 0 \end{pmatrix}, \; y = \frac{1-\sqrt{1-x}}{x\sqrt{1-x}}$$

Proof. Find the projection $J_I \in \mathcal{A}^+$ on the base $\mathcal{AO}^+_g$ of bundle $\mathcal{A}^+$, using results of [6]. The skew-symmetric form $\omega(X,Y) = \frac{1}{2}(g(J_IX,Y) - g(X,J_IY))$ corresponds to structure J_I. The matrix of form ω in the standard frame:

$$\omega = \frac{2}{\sqrt{-\tau}} \begin{pmatrix} 0 & B^* \\ -B^{*T} & 0 \end{pmatrix}$$

Corresponding skew-symmetric operator D, such that $\omega(X,Y) = g(DX,Y)$ is:

$$D = \frac{2}{\sqrt{-\tau}} \begin{pmatrix} 0 & -B^* \\ B^{*T} & 0 \end{pmatrix}$$

Then

$$(-D^2)^{-\frac{1}{2}} = \begin{pmatrix} (-\frac{4B^*B^{*T}}{\tau})^{-1/2} & 0 \\ 0 & (-\frac{4B^{*T}B^*}{\tau})^{-1/2} \end{pmatrix}$$

Find $(B^*B^{*T})^{-1/2}$ and $(B^{*T}B^*)^{-1/2}$. Use condition $J_I^2 = -1$, it is equivalent to system of equations:

$$\begin{cases} (1-2A^*)^2 - 4B^*B^{*T} = \tau E, \\ B^*C^* = A^*B^*, \\ (1-2C^*)^2 - 4B^{*T}B^* = \tau E. \end{cases}$$

Then $\left(-\frac{4B^*B^{*T}}{\tau}\right)^{-\frac{1}{2}} = \left(1 + \frac{(1-2A^*)^2}{-\tau}\right)^{-\frac{1}{2}}$ and $\left(-\frac{4B^{*T}B^*}{\tau}\right)^{-\frac{1}{2}} = \left(1 + \frac{(1-2C^*)^2}{-\tau}\right)^{-\frac{1}{2}}$.

By direct calculations one can show that $A^{*2} = xA^*$. Then

$$1 + \frac{(1-2A^*)^2}{-\tau} = 1 + \frac{(1+4(x-1)A^*)}{-\tau} = (1 + \frac{1}{-\tau})(1 - A^*).$$

$$\left(1+\frac{(1-2A^*)^2}{-\tau}\right)^{-\frac{1}{2}} = \frac{1}{\sqrt{1-\frac{1}{\tau}}}(1-A^*)^{-\frac{1}{2}} =$$

$$= \frac{1}{\sqrt{1-\frac{1}{\tau}}}(1+\frac{1}{2}A^*+\frac{3}{2^2}A^{*2}+\frac{15}{2^3}A^{*3}+\dots) =$$

$$= \frac{1}{\sqrt{1-\frac{1}{\tau}}}(1+\frac{1}{2}A^*+\frac{3}{2^2}xA^*+\frac{15}{2^3}x^2A^*+\dots) =$$

$$= \frac{1}{x\sqrt{1-\frac{1}{\tau}}}(x-A^*+A^*(1+\frac{1}{2}x+\frac{3}{2^2}x^2+\frac{15}{2^3}x^3+\dots)) =$$

$$= \frac{1}{x\sqrt{1-\frac{1}{\tau}}}(x+A^*(1-x)^{-\frac{1}{2}}-A^*) = \frac{1}{\sqrt{1-\frac{1}{\tau}}}\left(1+\frac{1-\sqrt{1-x}}{x\sqrt{1-x}}A^*\right).$$

Projection of almost complex structure J_I on $\mathcal{AO}_g^+$ is:

$$\pi(J_I) = (-D^2)^{-\frac{1}{2}}D = \frac{2}{\sqrt{1-\tau}}\begin{pmatrix} 0 & -(1+yA^*)B^* \\ (1+yC^*)B^{*T} & 0 \end{pmatrix},$$

therefore $\omega_J = \frac{2}{\sqrt{1-\tau}}\begin{pmatrix} 0 & (1+yA^*)B^* \\ -(1+yC^*)B^{*T} & 0 \end{pmatrix}$.

□

Remark 4. As $\pi(J_I) \in \mathcal{AO}_g^+$, then matrix $X = \frac{2}{\sqrt{1-\tau}}(1+yA^*)B^* \in SO(3)$. So the matrix of form ω is canonical $\begin{pmatrix} 0 & E \\ -E & 0 \end{pmatrix}$ in the frame $(u_1,u_2,u_3,u_4,u_5,u_6) = (e_1,e_2,e_3,Xe_4,Xe_5,Xe_6)$. The group $SO(3)\times SO(3)$ acts on $M = S^3\times S^3 = SU(2)\times SU(2)$ with canonical metric g by an isometry [2]. We have $[u_4,u_5]=u_6$, $[u_4,u_6]=-u_5$, $[u_5,u_6]=u_4$, $u_4,u_5,u_6 \in \{0\}\times\mathfrak{su}(2)$. Therefore frames (e) and (u) are equivalent [2]. One can repeat all the above calculations for the almost complex structure $I \in \mathcal{AO}_g^+$ in the new frame (u), and get $\pi(J_I) = I_0$, where $I_0 \in \mathcal{AO}_g^+$: $I_0u_4 = u_1$, $I_0u_5 = u_2$, $I_0u_6 = u_3$.

Lemma 3. *The form ω_J and metric g_I*

$$\omega_J = \begin{pmatrix} 0 & E \\ -E & 0 \end{pmatrix} \qquad g_I = \begin{pmatrix} 2t\left(1-\frac{A^*}{1+t}\right) & -1+2A^* \\ -1+2A^* & 2t\left(1-\frac{A^*}{1+t}\right) \end{pmatrix},$$

where $t=\sqrt{1-x}$ in the frame $(u_1,u_2,u_3,u_4,u_5,u_6)$.

Corollary. *Map*

$$I \in \mathcal{AO}_- \longrightarrow^{\alpha} J_I \longrightarrow^{\pi} \pi(J_I) \longrightarrow^{\alpha} J_{\pi(J_I)}$$

gives the almost complex structure, for which $(g_{\mathcal{NK}}, J_{\pi(J_I)}) \in \mathcal{NK}$.

Remark 5. As each almost complex structure J_I for $I \in \mathcal{AO}_-$ is in $\mathcal{A}^+_{\omega_J}$, then we have almost Hermitian structure (g_I, J_I), where $g_I(X,Y) = \omega_J(X, J_I Y)$ for all $I \in \mathcal{AO}_-$.

Theorem 3.*Ricci curvature of metric* g_I *is*

$$Ric_{11} = Ric_{22} = -\frac{8t^4 - 16t^3 - 10t^2 + 10t + 3}{2(4t^2-1)^2}; Ric_{33} = \frac{4t^2+1}{2(4t^2-1)^2};$$

$$Ric_{44} = Ric_{55} = -\frac{8t^4 - 16t^3 + 6t^2 - 2t - 1}{2(4t^2-1)^2}; Ric_{66} = \frac{-3 + 16t^4 - 8t^2}{2(4t^2-1)^2};$$

$$Ric_{ij} = 0, \ \ \textit{if } i \neq j$$

in the orthonormal frame for g_I. *The scalar curvature* $s = -\frac{8t^4 - 32t^3 - 2t^2 + 8t + 3}{(4t^2-1)^2}$.

Proof. In case $a_1^2 + a_3^2 \neq 0$ proper values and proper vectors of matrix g_I in the frame (u) are:

$$\begin{array}{lll}
\lambda_1 = 2t-1: & v_1 = \frac{1}{\sqrt{2(a_1^2+a_3^2)(2t-1)}}(-a_1, 0, a_3, -a_1, 0, a_3); \\
 & v_2 = \frac{1}{\sqrt{2(a_1^2+a_3^2)(1-t^2)(2t-1)}}(a_2a_3, a_1^2+a_3^2, a_1a_2, a_2a_3, a_1^2+a_3^2, a_1a_2); \\
\lambda_2 = 1: & v_3 = \frac{1}{\sqrt{2(1-t^2)}}(-a_3, a_2, -a_1, -a_3, a_2, -a_1); \\
\lambda_3 = 2t+1: & v_4 = \frac{1}{\sqrt{2(a_1^2+a_3^2)(2t+1)}}(-a_1, 0, a_3, a_1, 0, -a_3); \\
 & v_5 = \frac{1}{\sqrt{2(a_1^2+a_3^2)(2t+1)}}(a_2a_3, a_1^2+a_3^2, a_1a_2, -a_2a_3, -a_1^2-a_3^2, -a_1a_2); \\
\lambda_4 = 4t^2-1: & v_6 = \frac{1}{\sqrt{2(1-t^2)(4t^2-1)}}(-a_3, a_2, -a_1, a_3, -a_2, a_1)
\end{array}$$

In case $a_1^2 + a_3^2 = 0, \ a_2 \neq 0$ proper values and proper vectors of matrix g_I in the frame (u) are:

$$\begin{array}{lll}
\lambda_1 = 2t-1: & v_1 = \frac{1}{\sqrt{2(2t-1)}}(1, 0, 0, 1, 0, 0); \\
 & v_2 = \frac{1}{\sqrt{2(2t-1)}}(0, 0, 1, 0, 0, 1); \\
\lambda_2 = 1: & v_3 = \frac{1}{\sqrt{2(1-t^2)}}(0, a_2, 0, 0, a_2, 0); \\
\lambda_3 = 2t+1: & v_4 = \frac{1}{\sqrt{2(2t+1)}}(1, 0, 0, -1, 0, 0); \\
 & v_5 = \frac{1}{\sqrt{2(2t+1)}}(0, 0, 1, 0, 0, -1); \\
\lambda_4 = 4t^2-1: & v_6 = \frac{1}{\sqrt{2(1-t^2)(4t^2-1)}}(0, a_2, 0, 0, -a_2, 0)
\end{array}$$

In case $a_1^2 + a_2^2 + a_3^2 = 0$ we have $t = 1$ and

$$\begin{array}{rl}
\lambda_1 = 1: & v_1 = \frac{1}{\sqrt{2}}(1,0,0,1,0,0); \\
 & v_2 = \frac{1}{\sqrt{2}}(0,1,0,0,1,0); \\
 & v_3 = \frac{1}{\sqrt{2}}(0,0,1,0,0,1); \\
\lambda_2 = 3: & v_4 = \frac{1}{\sqrt{6}}(1,0,0,-1,0,0); \\
 & v_5 = \frac{1}{\sqrt{6}}(0,1,0,0,-1,0); \\
 & v_6 = \frac{1}{\sqrt{6}}(0,0,1,0,0,-1)
\end{array}$$

The direct calculations by formulas from [1] gives tensor Ricci in the frame $(v) = (v_1, v_2, v_3, v_4, v_5, v_6)$.

References

[1] Besse A.L. Einstein manifolds. Springer-Verlag Berlin, Heidelberg,1987.

[2] Butruille J.B. Homogeneous nearly Kähler manifolds, math.DG/0612655v1

[3] Hitchin N. Stable forms and special metrics,math.DG/0107101;

[4] Hitchin N. The geometry of three-forms in six and seven dimensions,math.DG/0010054v1;

[5] Kobayashi S.,Nomizu K.:"Foundations of differential geometry", Vol.2, *Interscienѕe Publishers*, New York, London, (1969).

[6] Daurtseva N.A. On the manifold of almost complex structures. Mat. Zametki, 2005, 78:1, 66-71